

# Tail estimates of series of independent random variables

KRZYSZTOF ZAJKOWSKI <sup>1</sup>

Institute of Mathematics, University of Białystok  
Akademicka 2, 15-267 Białystok, Poland  
kryza@math.uwb.edu.pl

## Abstract

Variational formulas for tail estimates (rate functions) of series of independent random variables are given.

*2010 Mathematics Subject Classification:* 44A15, 60F10

*Key words:* Cramér transform, Legendre-Fenchel transform, contraction principle (large deviations theory)

## 1 Introduction

For series  $\sum t_i g_i$  of independent standard normal r.v.'s, where  $\sum t_i^2 < \infty$  and  $g_i \in \mathcal{N}(0, 1)$ , it is well-known their tail estimations of the form

$$\Pr\left(\sum t_i g_i > \alpha\right) \leq \exp\left(-\frac{\alpha^2}{2\sum t_i^2}\right);$$

see for instance [6]. In Section 2 we present a derivation of this estimation and show some variational formula on the function  $\frac{\alpha^2}{2\sum t_i^2}$ . It will turn out that it is given by

$$\frac{\alpha^2}{2\sum t_i^2} = \inf_{\sum t_i b_i = \alpha} \sum b_i^2.$$

In Section 3 we prove variational formulas for a wider class of series of independent random variables. Let us stress that they mostly do not possess simple forms as in the above Gaussian case.

The function  $\frac{\alpha^2}{2\sum t_i^2}$  is the *Cramér transform* of the random series  $\sum t_i g_i$ . The Cramér transform is the *Legendre-Fenchel transform* of the *cumulant generating function* of r.v. To realize our purposes we will need the general notion of the Legendre-Fenchel transform in topological spaces (see [3] or [1]). Let  $V$  be a real locally convex Hausdorff space and  $V^*$  its dual space. By  $\langle \cdot, \cdot \rangle$  we denote the canonical pairing between  $V$  and  $V^*$ . Let  $f : V \mapsto \mathbb{R} \cup \{\infty\}$  be a function nonidentically  $\infty$ . By  $\mathcal{D}(f)$  we denote the

---

<sup>1</sup>The author is supported by the Polish National Science Center, Grant no. DEC-2011/01/B/ST1/03838.

*effective domain* of  $f$ , i.e.  $\mathcal{D}(f) = \{u \in V : f(u) < \infty\}$ . A function  $f^* : V^* \mapsto \mathbb{R} \cup \{\infty\}$  defined by

$$f^*(u^*) = \sup_{u \in V} \{\langle u, u^* \rangle - f(u)\} = \sup_{u \in \mathcal{D}(f)} \{\langle u, u^* \rangle - f(u)\} \quad (u^* \in V^*)$$

is called the *Legendre-Fenchel transform* (*convex conjugate*) of  $f$  and a function  $f^{**} : V \mapsto \mathbb{R} \cup \{\infty\}$  defined by

$$f^{**}(u) = \sup_{u^* \in V^*} \{\langle u, u^* \rangle - f^*(u^*)\} = \sup_{u^* \in \mathcal{D}(f^*)} \{\langle u, u^* \rangle - f^*(u^*)\} \quad (u \in V)$$

is called the *convex biconjugate* of  $f$ .

The functions  $f^*$  and  $f^{**}$  are convex and lower semicontinuous in the weak\* and weak topology on  $V^*$  and  $V$ , respectively. Moreover, the *biconjugate theorem* states that the function  $f : V \mapsto \mathbb{R} \cup \{\infty\}$  not identically equal to  $+\infty$  is convex and lower semicontinuous if and only if  $f = f^{**}$ .

Let us mention additional properties of the convex conjugates; see 4.3 Examples in [3]. Let  $V$  be a normed space. We denote by  $\|\cdot\|$  the norm of  $V$  and by  $\|\cdot\|_*$  the norm of  $V^*$ . For conjugates exponents  $p, q \in (1, \infty)$  ( $\frac{1}{p} + \frac{1}{q} = 1$ ), a function  $\frac{1}{q}\|u^*\|_*^q$  is the convex conjugate of  $\frac{1}{p}\|u\|^p$ .

**Remark 1.1.** Let us emphasize that in Hilbert spaces a function  $\frac{1}{2}\|u\|^2$  one can treat as the function invariant with respect to the Legendre-Fenchel transform.

Let us list another properties and one lemma. The convex-conjugation is order-reversing:

$$\text{if } f \leq g \text{ then } f^* \geq g^* \quad (1)$$

and

$$\text{if } g(u) = f(au) \text{ where } (a \neq 0) \text{ then } g^*(u^*) = f^*\left(\frac{u^*}{a}\right). \quad (2)$$

Let  $V$  and  $W$  be normed spaces. The dual space of their product  $(V \times W)^*$  is isomorphic to a direct sum of their dual spaces  $V^* \oplus W^*$  in this sense that a canonical pairing between  $V \times W$  and  $(V \times W)^*$  is given by the sum  $\langle u, u^* \rangle + \langle v, v^* \rangle$  where  $u \in V$ ,  $v \in W$ ,  $u^* \in V^*$  and  $v^* \in W^*$ .

**Lemma 1.2.** Let  $V$  and  $W$  be normed spaces, functions  $f : V \mapsto \mathbb{R} \cup \{\infty\}$  and  $g : W \mapsto \mathbb{R} \cup \{\infty\}$ . The convex conjugate of a function  $\varphi(u, v) = f(u) + g(v)$  defined on  $V \times W$  equals

$$\varphi^*(u^*, v^*) = f^*(u^*) + g^*(v^*)$$

*Proof.* The convex conjugate of  $\varphi$  equals

$$\begin{aligned}
\varphi^*(u^*, v^*) &= \sup_{(u, v) \in V \times W} \{ \langle u, u^* \rangle + \langle v, v^* \rangle - f(u) - g(v) \} \\
&= \sup_{u \in V} \sup_{v \in W} \{ \langle u, u^* \rangle - f(u) + \langle v, v^* \rangle - g(v) \} \\
&= \sup_{u \in V} \{ \langle u, u^* \rangle - f(u) \} + \sup_{v \in W} \{ \langle v, v^* \rangle - g(v) \} \\
&= f^*(u^*) + g^*(v^*).
\end{aligned}$$

□

## 2 A model example

The moment generating function of a standard normal r.v.  $g$  equals  $Ee^{sg} = e^{\frac{s^2}{2}}$  and its cumulant generating function

$$\psi_g(s) := \ln Ee^{sg} = \frac{s^2}{2}.$$

The function  $\frac{s^2}{2}$  is invariant with respect to Legendre transform (see Remark 1.1) that is the Cramér transform of  $g$  equals

$$\psi_g^*(\alpha) = \frac{\alpha^2}{2}.$$

Let  $I \subset \mathbb{N}$  and  $(g_i)_{i \in I}$  be a sequence of independent standard normal r.v.'s. For  $\mathbf{t} = (t_i)_{i \in I} \in \ell^2(I) \equiv \ell^2$  consider  $X_{\mathbf{t}} = \sum_{i \in I} t_i g_i$  (convergence in  $L^2$  and a.s.). Observe that  $EX_{\mathbf{t}}^2 = \sum_{i \in I} t_i^2 = \|\mathbf{t}\|^2$ . Thus a subset  $\{X_{\mathbf{t}} : \mathbf{t} \in \ell^2\}$  of  $L^2$  is isomorphic to  $\ell^2$ . The cumulant generating function of  $X_{\mathbf{t}}$  we will denote by  $\psi_{\mathbf{t}}$ , i.e.  $\psi_{\mathbf{t}} = \psi_{X_{\mathbf{t}}}$ . Notice that for fixed  $s$  we can consider  $\psi_{\mathbf{t}}(s)$  as a functional of the variable  $\mathbf{t}$  in  $\ell^2$ . We will denote it by  $\psi^s$ . Let us emphasize that  $\psi^s(\mathbf{t}) = \psi_{\mathbf{t}}(s)$ .

Let  $I_N$  denote a set  $\{1, 2, \dots, N\} \cap I$ . For  $\mathbf{t} \in \ell^2(I)$  let  $\mathbf{t}^N$  is defined as follows:  $t_i^N = t_i$  for  $i \in I_N$  and  $t_i^N = 0$  for  $i \in I \setminus I_N$ . Observe that  $\mathbf{t}^N \xrightarrow{\ell^2} \mathbf{t}$  and

$$\psi^s(\mathbf{t}^N) = \ln Ee^{s \sum_{i \in I_N} t_i g_i} = \sum_{i \in I_N} \psi_g(st_i) = \frac{s^2}{2} \sum_{i \in I_N} t_i^2 \rightarrow \frac{s^2}{2} \|\mathbf{t}\|^2$$

as  $N \rightarrow \infty$ . For each  $s \in \mathbb{R}$  the function  $\frac{s^2}{2} \|\mathbf{t}\|^2$  is continuous on  $\ell^2$ . It follows that  $\psi^s(\mathbf{t}) = \frac{s^2}{2} \|\mathbf{t}\|^2$ . The function  $\frac{1}{2} \|\mathbf{t}\|^2$  is invariant with respect to the convex conjugate on  $\ell^2$  (see Remark 1.1). Taking in (2)  $f(\mathbf{t}) = \frac{1}{2} \|\mathbf{t}\|^2$  and  $a = s$  we have  $\psi^s(\mathbf{t}) = f(s\mathbf{t})$  and get

$$(\psi^s)^*(\mathbf{a}) = f^*\left(\frac{\mathbf{a}}{s}\right) = \frac{1}{2} \left\| \frac{\mathbf{a}}{s} \right\|^2$$

for  $s \neq 0$ . The function  $\psi^s(\mathbf{t}) = \frac{s^2}{2}\|\mathbf{t}\|_2^2$  is convex and continuous on  $\ell^2$ . By the biconjugate theorem we have

$$\begin{aligned}\psi^s(\mathbf{t}) &= \sup_{\mathbf{a} \in \ell^2} \{ \langle \mathbf{t}, \mathbf{a} \rangle - (\psi^s)^*(\mathbf{a}) \} \\ &= \sup_{\mathbf{a} \in \ell^2} \left\{ \langle \mathbf{t}, \mathbf{a} \rangle - \frac{1}{2} \left\| \frac{\mathbf{a}}{s} \right\|^2 \right\}.\end{aligned}$$

Substituting  $\mathbf{a} = s\mathbf{b}$  into the right-hand side we may rewrite the above as follows

$$\psi^s(\mathbf{t}) = \sup_{\mathbf{b} \in \ell^2} \left\{ s \langle \mathbf{t}, \mathbf{b} \rangle - \frac{1}{2} \|\mathbf{b}\|^2 \right\}.$$

If we split the supremum into two parts: over  $\mathbb{R}$  and hyperplanes  $\{\mathbf{b} \in \ell^2 : \langle \mathbf{b}, \mathbf{t} \rangle = \text{constant}\}$  then we get

$$\begin{aligned}\psi_{\mathbf{t}}(s) &= \sup_{\alpha \in \mathbb{R}} \sup_{\substack{\mathbf{b} \in \ell^2: \\ \langle \mathbf{t}, \mathbf{b} \rangle = \alpha}} \left\{ s \langle \mathbf{t}, \mathbf{b} \rangle - \frac{1}{2} \|\mathbf{b}\|^2 \right\} \\ &= \sup_{\alpha \in \mathbb{R}} \left\{ s\alpha - \frac{1}{2} \inf_{\substack{\mathbf{b} \in \ell^2: \\ \langle \mathbf{t}, \mathbf{b} \rangle = \alpha}} \|\mathbf{b}\|^2 \right\}.\end{aligned}\tag{3}$$

We should remember that  $\psi^s(\mathbf{t}) = \psi_{\mathbf{t}}(s)$  that is  $\psi_{\mathbf{t}}(s) = \frac{1}{2}(\|\mathbf{t}\|s)^2$ . Using again (2) but now for  $f(x) = \frac{1}{2}x^2$  and  $a = \|\mathbf{t}\|$  we obtain that

$$\psi_{\mathbf{t}}^*(\alpha) = \frac{\alpha^2}{2\|\mathbf{t}\|^2}.$$

It is the above-mentioned tail estimate for the separable Gaussian process given by series  $\sum t_i g_i$ .

The function  $\psi_{\mathbf{t}}(s) = \frac{1}{2}(\|\mathbf{t}\|s)^2$  is convex and continuous on  $\mathbb{R}$ . By the biconjugate theorem we have that

$$\psi_{\mathbf{t}}(s) = \sup_{\alpha \in \mathbb{R}} \left\{ s\alpha - \frac{\alpha^2}{2\|\mathbf{t}\|^2} \right\}.\tag{4}$$

Comparing (3) and (4) we see (for a general case see Th.3.3) that

$$\frac{\alpha^2}{2\|\mathbf{t}\|^2} = \frac{1}{2} \inf_{\substack{\mathbf{b} \in \ell^2: \\ \langle \mathbf{t}, \mathbf{b} \rangle = \alpha}} \|\mathbf{b}\|^2,$$

what we could also check by using the Lagrange multipliers technique.

### 3 Main Theorem

The cumulant generating function  $\psi_X(s) = \ln Ee^{sX}$  of any random variable  $X$  is convex and lower semicontinuous on  $\mathbb{R}$  (analytic on  $\text{int}\mathcal{D}(\psi_X)$ ). It maps  $\mathbb{R}$  into  $\mathbb{R} \cup \{\infty\}$  and takes value zero at zero but it is possible that  $\psi_X(s) = \infty$  when  $s \neq 0$ . It is reasonable to assume that it is finite on some neighborhood of zero, i.e.  $X$  satisfies condition:  $\exists \lambda > 0$  s.t.  $Ee^{\lambda|X|} < \infty$ . Let us emphasize that if  $EX = 0$  then  $\psi_X \geq 0$  but the Cramér transform  $\psi_X^*$  is always nonnegative and attains 0 at the value  $EX$ .

Before proving our main theorem, we show forms of the cumulant generating function and the Cramér transform of series of independent random variables.

**Proposition 3.1.** *Let  $(X_i)_{i \in I}$  be a sequence of zero-mean independent random variables with common bounded second moments. Let for each  $i \in I$  the cumulant generating function  $\psi_i(s) := \psi_{X_i}(s) = \ln Ee^{sX_i}$  is finite on some neighborhood of zero. Then for each  $\mathbf{t} = (t_i)_{i \in I} \in \ell^2$  the cumulant generating function of a random series  $X_{\mathbf{t}} = \sum_{i \in I} t_i X_i$  is given by the following equality*

$$\psi_{\mathbf{t}}(s) := \psi_{X_{\mathbf{t}}}(s) = \sum_{i \in I} \psi_i(st_i).$$

*Proof.* Because  $(X_i)$  are independent, centered and have common bounded second moments then for every  $\mathbf{t} \in \ell^2$  the series  $X_{\mathbf{t}} = \sum_{i \in I} t_i X_i$  converges in  $L^2$  and a.s.. Let us emphasize that the convergence of series  $X_{\mathbf{t}}$  in  $L^2$  is equivalent to the convergence of sequences  $\mathbf{t}$  in  $\ell^2$ . By fixed  $s$  we can consider the cumulant generating function  $\psi_{\mathbf{t}}(s)$  as a functional of the variable  $\mathbf{t}$  in  $\ell^2$ . We will denote it by  $\psi^s(\mathbf{t})$ , i.e.  $\psi^s(\mathbf{t}) = \psi_{\mathbf{t}}(s)$ . We show that for every  $s \in \mathbb{R}$  the functional  $\psi^s$  is convex and lower semicontinuous on  $\ell^2$ .

Convexity one may check by using the Hölder inequality. Let  $\mathbf{t}, \mathbf{u} \in \ell^2$  and  $\lambda \in (0, 1)$  then

$$\begin{aligned} \psi^s(\lambda \mathbf{t} + (1 - \lambda) \mathbf{u}) &= \ln Ee^{s \sum_{i \in I} (\lambda t_i + (1 - \lambda) u_i) X_i} \\ &= \ln E[(e^{s \sum_{i \in I} t_i X_i})^\lambda (e^{s \sum_{i \in I} u_i X_i})^{1 - \lambda}] = \ln E[(e^{sX_{\mathbf{t}}})^\lambda (e^{sX_{\mathbf{u}}})^{1 - \lambda}]. \end{aligned}$$

By the Hölder inequality for exponents  $1/\lambda$  and  $1/(1 - \lambda)$  we get

$$E[(e^{sX_{\mathbf{t}}})^\lambda (e^{sX_{\mathbf{u}}})^{1 - \lambda}] \leq (Ee^{sX_{\mathbf{t}}})^\lambda (Ee^{sX_{\mathbf{u}}})^{1 - \lambda}$$

and, in consequence,

$$\begin{aligned} \psi^s(\lambda \mathbf{t} + (1 - \lambda) \mathbf{u}) &\leq \lambda \ln Ee^{sX_{\mathbf{t}}} + (1 - \lambda) \ln Ee^{sX_{\mathbf{u}}} \\ &= \lambda \psi^s(\mathbf{t}) + (1 - \lambda) \psi^s(\mathbf{u}). \end{aligned}$$

Lower semicontinuity follows from Fatou's lemma. Let  $\mathbf{t}^n \rightarrow \mathbf{t}^0$  in  $\ell^2$ . Note that  $X_{\mathbf{t}^n}$  converges a.s. to  $X_{\mathbf{t}^0}$ . Then

$$\begin{aligned} \liminf_{n \rightarrow \infty} \psi^s(\mathbf{t}^n) &= \liminf_{n \rightarrow \infty} \ln E e^{s X_{\mathbf{t}^n}} \geq \ln E(\liminf_{n \rightarrow \infty} e^{s X_{\mathbf{t}^n}}) \\ &= \ln E(e^{s \lim_{n \rightarrow \infty} X_{\mathbf{t}^n}}) = \psi^s(\mathbf{t}^0). \end{aligned}$$

It means that  $\psi^s$  is lower semicontinuous on  $\ell^2$ .

Let  $\ell_0$  denote the space of sequences with finite supports. Observe that  $\ell_0$  is a dense subset of  $\ell^2$ . For  $\mathbf{t} \in \ell_0$  we have

$$\begin{aligned} \psi^s(\mathbf{t}) &= \ln E e^{s X_{\mathbf{t}}} = \ln \prod_{i \in I} e^{s t_i X_i} \\ &= \sum_{i \in I} \psi_i(s t_i). \end{aligned}$$

For  $\mathbf{t} \in \ell^2$  consider a series  $\sum_{i \in I} \psi_i(s t_i)$ . Since  $E X_i = 0$ ,  $\psi_i \geq 0$ . It follows that  $\sum_{i \in I} \psi_i(s t_i)$  is convergent or divergent to plus infinity. Since  $\psi_i$  are convex, this series defines a convex function on the whole  $\ell^2$ . Let  $\mathbf{t}^n \rightarrow \mathbf{t}^0$  in  $\ell^2$ . Hence for every  $i \in I$   $t_i^n \rightarrow t_i^0$ . By superadditivity of the limit inferior and, next, by lower semicontinuity of each  $\psi_i$ , we get

$$\begin{aligned} \liminf_{n \rightarrow \infty} \psi^s(\mathbf{t}^n) &= \liminf_{n \rightarrow \infty} \sum_{i \in I} \psi_i(s t_i^n) \geq \sum_{i \in I} \liminf_{n \rightarrow \infty} \psi_i(s t_i^n) \\ &\geq \sum_{i \in I} \psi_i(s t_i^0) = \psi^s(\mathbf{t}^0). \end{aligned}$$

Notice that both functions:  $\psi^s(\mathbf{t})$  and the series  $\sum_{i \in I} \psi_i(s t_i)$  are convex and lower semicontinuous on  $\ell^2$  and moreover coincide on  $\ell_0$  (a dense subset of  $\ell^2$ ). It follows that these functions are equal on whole  $\ell^2$ , i.e.

$$\psi^s(\mathbf{t}) = \sum_{i \in I} \psi_i(s t_i)$$

for every  $\mathbf{t}$  in  $\ell^2$ . □

Let us observe that for  $s = 0$   $\psi^0 \equiv 0$  and its convex conjugate  $(\psi^0)^*(\mathbf{a}) = 0$  for  $\mathbf{a} = \mathbf{0}$  and  $\infty$  otherwise. From now on we assume that  $s \neq 0$ . A form of  $(\psi^s)^*$  for  $s \neq 0$  is described in the following:

**Proposition 3.2.** *Under the assumptions of Proposition 3.1. The convex conjugate of  $\psi^s(\mathbf{t}) = \sum_{i \in I} \psi_i(s t_i)$  defined on  $\ell^2$  equals*

$$(\psi^s)^*(\mathbf{a}) = \sum_{i \in I} \psi_i^*\left(\frac{a_i}{s}\right) \quad (s \neq 0)$$

for  $\mathbf{a} \in \ell^2$ , where  $\psi_i^*$ 's are the Cramér transforms of  $X_i$ 's.

*Proof.* The convex conjugate  $(\psi^s)^*$  is convex and lower semicontinuous on  $(\ell^2)^* \simeq \ell^2$ . Assume first that  $I$  is a finite set. By virtue of the form of  $\psi^s$ , Lemma 1.2 and the property (2), for  $\mathbf{a}$  in  $\ell^2(I)$ , we get

$$(\psi^s)^*(\mathbf{a}) = \sum_{i \in I} \psi_i^*\left(\frac{a_i}{s}\right) \quad (s \neq 0).$$

Define now a functional  $\sum_{i \in I} \psi_i^*\left(\frac{a_i}{s}\right)$  on whole space  $\ell^2$ . Since  $\psi_i^*$ 's are convex and lower semicontinuous, this functional is convex and, similarly as in the case of  $\sum_{i \in I} \psi_i$ , one can show that it is also lower semicontinuous on  $\ell^2$ . Because this functional coincides with  $(\psi^s)^*$  on the dense subspace  $\ell_0$  then both functionals are equal on  $\ell^2$ .  $\square$

Let us emphasize that the functions  $(\psi^s)^*$  are nonnegative and lower semicontinuous. In large deviation theory such functions are called rate functions (good rate functions when level sets are not only closed but also compact). In the main theorem below we show that the contraction principle applied to the function  $(\psi^1)^*$  by using a functional  $\langle \mathbf{t}, \cdot \rangle$  over  $\ell^2$  gives the Cramér transform of  $X_{\mathbf{t}}$ .

**Theorem 3.3.** *Let a sequence of r.v.'s  $(X_i)_{i \in I}$  satisfies the assumptions of Proposition 3.1. Then for every  $\mathbf{t} = (t_i)_{i \in I} \in \ell^2$  the Cramér transform  $\psi_{X_{\mathbf{t}}}^* = \psi_{\mathbf{t}}^*$  of a random series  $X_{\mathbf{t}} = \sum_{i \in I} t_i X_i$  is given by the following variational formula*

$$\psi_{\mathbf{t}}^*(\alpha) = \inf_{\mathbf{b} \in \ell^2: \langle \mathbf{t}, \mathbf{b} \rangle = \alpha} \sum_{i \in I} \psi_i^*(b_i),$$

for  $\alpha \in \text{int}\mathcal{D}(\psi_{\mathbf{t}}^*)$ , where  $\psi_i^*$ 's are the Cramér transform of  $X_i$ 's.

*Proof.* The functional  $\psi^s$  is convex and lower semicontinuous on  $\ell^2$ . By virtue of the biconjugate theorem we have

$$\psi^s(\mathbf{t}) = \sup_{\mathbf{a} \in \ell^2} \{ \langle \mathbf{t}, \mathbf{a} \rangle - (\psi^s)^*(\mathbf{a}) \},$$

where  $(\psi^s)^*(\mathbf{a}) = \sum_{i \in I} (\psi_i)^*\left(\frac{a_i}{s}\right)$  ( $s \neq 0$ ). Substituting  $\mathbf{a} = s\mathbf{b}$  we get

$$(\psi^s)^*(\mathbf{a}) = \sum_{i \in I} \psi_i^*\left(\frac{a_i}{s}\right) = \sum_{i \in I} \psi_i^*(b_i) = (\psi^1)^*(\mathbf{b})$$

and we can rewrite the above as follows

$$\psi^s(\mathbf{t}) = \sup_{\mathbf{b} \in \ell^2} \{ s \langle \mathbf{t}, \mathbf{b} \rangle - (\psi^1)^*(\mathbf{b}) \}. \quad (5)$$

Let us return to the function  $\psi_{\mathbf{t}}$  which is convex and lower semicontinuous on  $\mathbb{R}$ . By the biconjugate theorem we have

$$\psi_{\mathbf{t}}(s) = \sup_{\alpha \in \mathbb{R}} \{ s\alpha - \psi_{\mathbf{t}}^*(\alpha) \}.$$

Let us recall that  $\psi_{\mathbf{t}}(s) = \psi^s(\mathbf{t})$ . If we split the supremum of (5), as in Model Example, into two parts then we get

$$\begin{aligned}\psi_{\mathbf{t}}(s) &= \psi^s(\mathbf{t}) = \sup_{\alpha \in \mathbb{R}} \sup_{\substack{\mathbf{b} \in \ell^2: \\ \langle \mathbf{t}, \mathbf{b} \rangle = \alpha}} \{s \langle \mathbf{t}, \mathbf{b} \rangle - (\psi^1)^*(\mathbf{b})\} \\ &= \sup_{\alpha \in \mathbb{R}} \{s\alpha - \inf_{\mathbf{b} \in \ell^2: \langle \mathbf{t}, \mathbf{b} \rangle = \alpha} (\psi^1)^*(\mathbf{b})\}.\end{aligned}$$

Let  $\varphi_{\mathbf{t}}(\alpha)$  denote the function  $\inf_{\mathbf{b} \in \ell^2: \langle \mathbf{t}, \mathbf{b} \rangle = \alpha} (\psi^1)^*(\mathbf{b})$ . The functional  $(\psi^1)^*$  is convex on  $\ell^2$ . Convexity is preserved under contraction by linear transformation (see [4, Th.III.32]). It suffices to state that  $\varphi_{\mathbf{t}}$  and  $\psi_{\mathbf{t}}^*$  coincide on  $\text{int}\mathcal{D}(\psi_{\mathbf{t}}^*)$  (both ones take  $\infty$  on a complement of  $\text{cl}\mathcal{D}(\psi_{\mathbf{t}}^*)$ ) that is

$$\psi_{\mathbf{t}}^*(\alpha) = \inf_{\mathbf{b} \in \ell^2: \langle \mathbf{t}, \mathbf{b} \rangle = \alpha} (\psi^1)^*(\mathbf{b}),$$

for  $\alpha \in \text{int}\mathcal{D}(\psi_{\mathbf{t}}^*)$ , where  $(\psi^1)^*(\mathbf{b}) = \sum_{i \in I} \psi_i^*(b_i)$ . □

**Remark 3.4.** We can not prove in general that  $\varphi_{\mathbf{t}}$  is lower semicontinuous. Sometimes it is obvious when for instance the effective domain of  $\psi_{\mathbf{t}}^*$  is open subset of  $\mathbb{R}$  (or even whole  $\mathbb{R}$ ; see Model Example and Example 3.5). Under an assumption that  $(\psi^1)^*$  is a good rate function with respect to weak\* topology we can state lower-semicontinuity of  $\varphi_{\mathbf{t}}$  (see Example 3.6).

**Example 3.5.** Let  $X$  be r.v. with a Laplace density  $\frac{1}{2}e^{-|x|}$ . Its moment generating function  $Ee^{sX} = \frac{1}{1-s^2}$  for  $|s| < 1$  and  $\infty$  otherwise. Let us observe that

$$Ee^{sX} = \frac{1}{1-s^2} = \sum_{n=0}^{\infty} s^{2n} \geq \sum_{n=0}^{\infty} \frac{s^{2n}}{n!2^n} = e^{\frac{s^2}{2}} = Ee^{sg},$$

where  $g$  is standard normal distributed. Thus  $\psi_X \geq \psi_g$  and, since the Cramér transform is order-reversing, we have

$$\psi_X^*(\alpha) \leq \psi_g^*(\alpha) = \frac{\alpha^2}{2}. \quad (6)$$

Moreover by using the classical Legendre transform we can calculate an evident form of  $\psi_X^*$  and get

$$\psi_X^*(\alpha) = \frac{\alpha^2}{\sqrt{1+\alpha^2}+1} + \ln \frac{2}{\sqrt{1+\alpha^2}+1}.$$

Consider a sequence  $(X_i)_{i \in I}$  of independent r.v.'s with the same Laplace distribution. By (6) we have

$$(\psi^1)^*(\mathbf{b}) = \sum_{i \in I} \psi_X^*(b_i) \leq \frac{1}{2} \|\mathbf{b}\|^2.$$

It means that  $(\psi^1)^*$  takes finite values on the whole space  $\ell^2$  and for every  $\alpha \in \mathbb{R}$  there is a finite infimum:

$$\inf_{\mathbf{b} \in \ell^2: \langle \mathbf{t}, \mathbf{b} \rangle = \alpha} \sum_{i \in I} \psi_X^*(b_i)$$

that is the finite value of  $\psi_{\mathbf{t}}^*$  at  $\alpha$ .

In the paper [7] one can find an example of variational formulas for the Cramér transform of series of weighted symmetric Bernoulli random variables but with coefficient belonging to the space  $\ell^1$ . In the context of our Theorem 3.3 we recall main result of this paper but now with coefficients in the bigger space  $\ell^2$ .

**Example 3.6.** If  $X$  is a symmetric Bernoulli r.v., i.e.  $Pr(X = \pm 1) = \frac{1}{2}$ , then  $Ee^{sX} = \cosh s$ . By power series expansions one has  $\cosh s \leq e^{\frac{s^2}{2}}$ . In this example, conversely as in previous one, we get that  $\psi_X(s) \leq \psi_g(s) = \frac{s^2}{2}$  and

$$\psi_X^*(\alpha) \geq \frac{\alpha^2}{2}. \quad (7)$$

One can check that

$$\psi_X^*(\alpha) = \frac{1}{2}[(1 + \alpha) \ln(1 + \alpha) + (1 - \alpha) \ln(1 - \alpha)]$$

for  $|\alpha| \leq 1$  and  $\infty$  otherwise; we take  $0 \ln 0 = 0$ . Note that  $\psi_X^*(\pm 1) = \ln 2$ .

For a sequence of independent Bernoulli r.v.'s, by the above inequality, we have

$$(\psi^1)^*(\mathbf{b}) \geq \frac{1}{2} \|\mathbf{b}\|^2.$$

Since  $(\psi^1)^*$  is lower semicontinuous in the weak\* topology, level sets  $\{\mathbf{b} \in \ell^2 : (\psi^1)^*(\mathbf{b}) \leq c\}$  are weak\* closed; for  $c < 0$  are empty sets. By the above, for each  $c \geq 0$  the level set of  $(\psi^1)^*$  is contained in a closed ball  $\overline{B}(\mathbf{0}; \sqrt{2c}) = \{\mathbf{b} \in \ell^2 : \|\mathbf{b}\| \leq \sqrt{2c}\}$ . By virtue of the Banach-Alaoglo theorem (see [5, Th. 3.15]) balls  $\overline{B}(\mathbf{0}; \sqrt{2c})$  are weak\* compact (are polar sets of balls  $\overline{B}(\mathbf{0}; 1/\sqrt{2c})$ ). It follows that the level sets  $\{\mathbf{b} \in \ell^2 : (\psi^1)^*(\mathbf{b}) \leq c\}$  are weak\* compact as closed subsets of compact sets. It means that in this topology  $(\psi^1)^*$  is a good rate function and by the contraction principle (see [2, Th.4.2.1])  $\varphi_{\mathbf{t}}$  is also good rate function on  $\mathbb{R}$ , in particular, it is lower semicontinuous.

Let us emphasize that if we know that  $(\psi^1)^*$  is a good rate function then we can prove lower-semicontinuity of  $\varphi_{\mathbf{t}}$ .

**Remark 3.7.** Eventually, let us emphasize once again that only in Gaussian cases we can solve the variational formula on exponential rate functions for tail estimates of series of independent r.v.'s. For non-Gaussian processes it remains at us only these variational principles.

## References

- [1] V. Barbu, T. Precupanu, *Convexity and Optimization in Banach Spaces*, 4th ed., Springer Monographs in Mathematics, Springer, Dordrecht, (2012)
- [2] A. Dembo, O. Zeitouni, *Large Deviations Techniques and Applications*. Corrected reprints of the second (1998) edition, Stochastic Modeling and Applied Probability, 38, Springer-Verlag, Berlin, (2010)
- [3] I. Ekeland, R. Témam, *Convex Analysis and Variational Problems*, Translated from French. Corrected reprint of the 1976 English edition. Classics in Applied Mathematics, 28, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, (1999)
- [4] F. den Hollander, *Large Deviations*, Fields Institute Monographs, 14, Providence, RI: American Mathematical Society (2000)
- [5] W. Rudin, *Functional Analysis*, 2nd ed., International Series in Pure and Applied Mathematics (1991)
- [6] M. Talagrand, *Majorizing measures: the generic chaining*, Ann. Probab. 24, 1049-1103 (1996)
- [7] K. Zajkowski, *Cramér transform of Rademacher series*, Positivity (2014), doi: 10.1007/s11117-014-0313-5.